



Popular Matchings in Complete Graphs

Ágnes Cseh^{1,2} · Telikepalli Kavitha³

Received: 9 April 2019 / Accepted: 1 December 2020 / Published online: 25 January 2021
© The Author(s) 2021

Abstract

Our input is a complete graph G on n vertices where each vertex has a strict ranking of all other vertices in G . The goal is to construct a matching in G that is *popular*. A matching M is popular if M does not lose a head-to-head election against any matching M' : here each vertex casts a vote for the matching in $\{M, M'\}$ in which it gets a better assignment. Popular matchings need not exist in the given instance G and the popular matching problem is to decide whether one exists or not. The popular matching problem in G is easy to solve for odd n . Surprisingly, the problem becomes NP-complete for even n , as we show here. This is one of the few graph theoretic problems efficiently solvable when n has one parity and NP-complete when n has the other parity.

Keywords Popular matching · Complexity · Stable matching

A preliminary version of this work appeared in FSTTCS 2018 [9].

Ágnes Cseh: Supported by the Federal Ministry of Education and Research of Germany (BMBF) in the KI-LAB-ITSE framework—project number 01IS19066, the Hungarian Academy of Sciences under its Momentum Programme (LP2016-3/2020), OTKA Grant K128611, and COST Action CA16228 European Network for Game Theory. Telikepalli Kavitha: This work was done while visiting the Hungarian Academy of Sciences, Budapest.

✉ Ágnes Cseh
agnes.cseh@hpi.de
Telikepalli Kavitha
kavitha@tifr.res.in

¹ Centre for Economic and Regional Studies, Institute of Economics, Budapest, Hungary

² Hasso Plattner Institute, University of Potsdam, Potsdam, Germany

³ Tata Institute of Fundamental Research, Mumbai, India

1 Introduction

Consider a complete graph $G = (V, E)$ on n vertices where each vertex ranks all other vertices in a strict order of preference. Such a graph is called a *roommates instance with complete preferences*. The problem of computing a stable matching in G is classical and well-studied. Recall that a matching M is stable if there is no *blocking pair* with respect to M , i.e., a pair (u, v) where both u and v prefer each other to their respective assignments in M .

Stable matchings need not always exist in a roommates instance. For example, the instance given in Fig. 1 on 4 vertices d_0, d_1, d_2, d_3 has no stable matching. (Here d_0 's top choice is d_1 , second choice is d_2 , and last choice is d_3 , and similarly for the other vertices.)

Irving [20] gave an efficient algorithm to decide if G admits a stable matching or not. In this paper we consider a notion that is more relaxed than stability: this is the notion of *popularity*. For any vertex u , a ranking over neighbors can be extended naturally to a ranking over matchings as follows: u prefers matching M to matching M' if (1) u is matched in M and unmatched in M' or (2) u is matched in both and it prefers its partner in M to its partner in M' . For any two matchings M and M' , let $\phi(M, M')$ be the number of vertices that prefer M to M' .

Definition 1 Let M be any matching in G . M is *popular* if $\phi(M, M') \geq \phi(M', M)$ for every matching M' in G .

Suppose an election is held between M and M' where each vertex casts a vote for the matching that it prefers. So $\phi(M, M')$ (similarly, $\phi(M', M)$) is the number of votes for M (resp., M'). A popular matching M never loses an election to another matching M' since $\phi(M, M') \geq \phi(M', M)$: thus it is a weak *Condorcet winner* [5, 6] in the corresponding voting instance. So popularity captures collective decision by the vertex set and it can be considered as a natural relaxation of stability.

The notion of popularity was first introduced in bipartite graphs in 1975 by Gärdenfors—popular matchings always exist in bipartite graphs since stable




Fig. 1 An instance that admits two popular matchings—marked by dotted blue and dashed orange edges—but no stable matching. The preference list of each vertex is illustrated by the numbers on its edges: a lower number indicates a more preferred neighbor (Color figure online)

matchings always exist here [13] and every stable matching is popular [14]. The proof that every stable matching is popular holds in non-bipartite graphs as well [4]; in fact, it is easy to show that every stable matching is a min-size popular matching [17]. Relaxing the constraint of stability to popularity allows us to find feasible matchings that may exist in instances that do not admit stable matchings; moreover, even when stable matchings exist, there may be popular matchings that achieve more “social good” (such as larger size), which might be relevant in many applications.

Observe that the instance in Fig. 1 has two popular matchings: $M_1 = \{(d_0, d_1), (d_2, d_3)\}$ and $M_2 = \{(d_0, d_2), (d_1, d_3)\}$. However as was the case with stable matchings, popular matchings also need not always exist in the given instance G . Just take, for example, the same instance as in Fig. 1, but without vertex d_0 . A complete graph G on an even number of vertices that has no popular matching is also easy to describe: take *two* copies of this instance on 3 vertices, i.e., d_1, d_2, d_3 with preferences as given in Fig. 1 (without vertex d_0) and three more vertices d'_1, d'_2, d'_3 whose preferences are analogous to d_1, d_2, d_3 , respectively. Since the instance has to be complete, add d_1, d_2, d_3 (similarly, d'_1, d'_2, d'_3) at the tail of preference lists of d'_1, d'_2, d'_3 (resp., d_1, d_2, d_3) in some arbitrary order. It is easy to check that this instance on 6 vertices has no popular matching.

The *popular roommates problem* is to decide if G admits a popular matching or not. When the graph is not complete, it is known that the popular roommates problem is NP-complete [11, 15]. Here we are interested in the complexity of the popular roommates problem when the input instance is complete.

Interestingly, several popular matching problems that are intractable in bipartite graphs become tractable in *complete bipartite* graphs. The min-cost popular matching problem in bipartite graphs is such a problem—this is NP-hard in a bipartite graph with incomplete lists [11], however it can be solved in polynomial time in a bipartite graph with complete lists [8]. The difference is due to the fact that while there is no compact extended formulation of the convex hull of edge incidence vectors of all popular matchings in a general bipartite graph [10], this polytope has a compact extended formulation in a complete bipartite graph.

It is a simple observation (see Sect. 2) that when n is *odd*, a matching in a complete graph G on n vertices is popular only if it is stable. Since there is an efficient algorithm to decide if G admits a stable matching or not, the popular roommates problem in a complete graph G can be efficiently solved when n is odd. We show the following result here.

Theorem 1 *Let G be a complete graph on n vertices, where n is even. The problem of deciding whether G admits a popular matching or not is NP-complete.*

So the popular roommates problem with complete preference lists is NP-complete for even n while it is easy to solve for odd n . Some problems possess an inherently different nature depending on the parity of some characteristic input parameter, such as Latin squares [21] or various problems in voting [12]. Popular matchings do not belong to this set of problems—note that the popular roommates problem is non-trivial for every $n \geq 5$, i.e., there are both “yes instances”

and “no instances” of size n . It is rare and unusual for a natural decision problem in combinatorial optimization to be efficiently solvable when n has one parity and become NP-complete when n has the other parity. We are not aware of any natural optimization problem on graphs that is non-trivially tractable when the cardinality of the vertex set has one parity, which becomes intractable for the other parity.

1.1 Background and related work

The first polynomial time algorithm for the stable roommates problem was given by Irving [20] in 1985. Roommates instances that admit stable matchings were characterized in [30]. New polynomial time algorithms for the stable roommates problem were given in [29, 31].

Algorithmic questions for popular matchings in bipartite graphs have been well-studied in the last decade [1, 8, 17, 19, 22–24]. Not much was known on popular matchings in non-bipartite graphs. Biró et al. [1] proved that validating whether a given matching is popular can be done in polynomial time, even when ties are present in the preference lists. It was shown in [18] that every roommates instance on n vertices admits a matching with *unpopularity factor* $O(\log n)$ and that it is NP-hard to compute a least unpopularity factor matching. It was shown in [19] that computing a max-weight popular matching in a roommates instance with edge weights is NP-hard, and more recently, that computing a max-size popular matching in a roommates instance is NP-hard [3].

The complexity of the popular roommates problem was open for several years [1, 7, 18, 19, 27] and two independent NP-completeness proofs [11, 15] of this problem were shown very recently. Interestingly, both these hardness proofs need “incomplete preference lists”, i.e., the underlying graph is *not* complete. The reduction in [15] is from a variant of the vertex cover problem called the *partitioned vertex cover* problem and we discuss the reduction in [11] in Sect. 1.2 below. So the complexity status of the popular roommates problem in a complete graph was an open problem and we resolve it here.

An interpretation of roommate instances with complete preference lists might be that each vertex finds every other vertex acceptable, or that being matched to any vertex is better than being unmatched, or that there is no outside option and the agents are all obliged to be matched within the market. Computational hardness for instances with complete lists has been investigated in various matching problems under preferences. An example is the three-sided stable matching problem with cyclic preferences: this involves three groups of participants, say, men, women, and dogs, where dogs have strictly ordered preferences over men only, men have preferences over women only, and finally, women only list the dogs. If these preferences are allowed to be incomplete, the problem of finding a *weakly* stable matching is known to be NP-complete [2]. Until very recently, it had been one of the most intriguing open questions in stable matchings [27, 32] as to whether the same problem becomes tractable when lists are complete. Lam and Plaxton gave a hardness proof for complete preference lists very recently, disproving the published conjectures [26].

1.2 Techniques

The 1-in-3 SAT problem is a well-known NP-complete problem [28]: it consists of a Boolean formula B in CNF where every clause has 3 literals (none negated) and the problem is to find a satisfying truth assignment to the variables in B such that every clause has exactly one literal set to true. We show a polynomial time reduction from 1-in-3 SAT to the popular roommates problem with complete lists.

Our construction is based on the reduction in [11] that proved the NP-completeness of the popular roommates problem. However there are several differences between our reduction and the reduction in [11]. The reduction in [11] considered a popular matching problem in bipartite graphs called the “exclusive popular set” problem and showed it to be NP-complete—when preference lists are complete, this problem can be easily solved. Thus the reduction in [11] needs incomplete preference lists.

The exclusive popular set problem asks if there is a popular matching in the given bipartite graph where the set of matched vertices is S , for a given even-sized subset S . A key step in the reduction in [11] from this problem in bipartite graphs to the popular matching problem in non-bipartite graphs merges all vertices outside S into a single node. Thus the total number of vertices in the non-bipartite graph used in [11] is *odd*. Moreover, the fact that popular matchings always exist in bipartite graphs is crucially used in this reduction. However in our setting, the whole problem is to decide if *any* popular matching exists in the given graph—thus there are no popular matchings that “always exist” here.

The reduction in [11] primarily uses the LP framework of popular matchings in bipartite graphs from [22, 23, 25] to analyze the structure of popular matchings in their instance. The LP framework characterizing popular matchings in non-bipartite graphs is more complex [25], so we use the combinatorial characterization of popular matchings [17] in terms of forbidden alternating paths/cycles to show that any popular matching in our instance will yield a 1-in-3 satisfying truth assignment for B . To show the converse, we use a dual certificate similar to the one used in [11] to prove the popularity of the matching that we construct using a 1-in-3 satisfying truth assignment for B .

Organization of the paper We discuss preliminaries in Section 2. Section 3 describes the construction of our complete graph G corresponding to a given 1-in-3 SAT formula B . Section 4 studies the structure of the graph G and Section 5 shows that any popular matching in G yields a 1-in-3 satisfying truth assignment for B . Section 6 completes the reduction by showing how to obtain a popular matching in G from any 1-in-3 satisfying truth assignment for B .

2 Preliminaries

This section contains a characterization of popular matchings from [17]. We also include a simple proof of the claim stated in Section 1 that when n is odd, every popular matching in G has to be stable.

Let M be any matching in $G = (V, E)$. For any pair $(u, v) \notin M$, define $\text{vote}_u(v, M)$ as follows: (here $M(u)$ is u 's partner in M and $M(u) = \text{null}$ if u is unmatched in M)

$$\text{vote}_u(v, M) = \begin{cases} + & \text{if } u \text{ prefers } v \text{ to } M(u); \\ - & \text{if } u \text{ prefers } M(u) \text{ to } v. \end{cases}$$

Label every edge (u, v) that does not belong to M by the pair $(\text{vote}_u(v, M), \text{vote}_v(u, M))$. Thus every non-matching edge has a label in $\{(\pm, \pm)\}$. For example, if we consider the matching marked by the dashed orange edges in Fig. 1, then (d_1, d_2) is labeled $(+, +)$, (d_2, d_3) is labeled $(+, -)$, (d_0, d_1) is labeled $(+, -)$, and (d_0, d_3) is labeled $(-, -)$. Note that an edge is labeled $(+, +)$ if and only if it is a blocking edge to M .

We remind the reader that an alternating path/cycle with respect to M is a path/cycle whose alternate edges belong to M : thus edges in this path/cycle alternate between belonging to the matching M and *not* belonging to the matching M . Let G_M be the subgraph of G obtained by deleting edges labeled $(-, -)$ from G . The following theorem characterizes popular matchings in G .

Theorem 2 ([17]) *M is popular in G if and only if G_M does not contain any of the following with respect to M :*

- (1) *an alternating cycle with a $(+, +)$ edge;*
- (2) *an alternating path with two distinct $(+, +)$ edges;*
- (3) *an alternating path with a $(+, +)$ edge and an unmatched vertex as an endpoint of the path.*

Using the above characterization, it can be easily checked whether a given matching is popular or not [17]. Thus our NP-hardness result implies that the popular roommates problem with complete preferences is NP-complete.

When n is odd. Recall the claim made in Sect. 1 that when n is odd, every popular matching in G has to be stable. A simple proof of this statement is included below.

Observation 1 ([16]) *Let G be a complete graph on n vertices, where n is odd. Any popular matching in G has to be stable.*

Proof Since n is odd and G is complete, any popular matching leaves exactly one vertex unmatched. Let M be a popular matching and let v be the vertex left unmatched in M . Consider a vertex u adjacent to v . We know that $(u, w) \in M$ for some $w \in V \setminus \{v\}$, and due to Part (3) in Theorem 2, no $(+, +)$ edge is incident to w . Since v is adjacent not only to u , but to all vertices in the graph, this holds for all $w \in V$, i.e., there is no $(+, +)$ edge incident to any vertex. Thus M is stable. \square

3 The graph G

Recall that B is the input formula to 1-in-3 SAT. We assume that B has κ variables X_1, \dots, X_κ . The graph G that we construct here consists of gadgets in 4 levels along with 2 special gadgets that we will call the D -gadget and Z -gadget. Gadgets in level 1 correspond to variables in the formula B while gadgets in levels 0, 2, and 3 correspond to clauses in B . Variants of the gadgets in levels 0–3 and the D -gadget were used in [11] while the Z -gadget is new.

An overview We will first show that any popular matching M in G uses only *intra-gadget* edges. Every gadget will be in one of the following two states in M : *stable* state or *unstable* state. A gadget is in stable state if and only if there is no edge e with both its endpoints in this gadget such that e blocks M . Our aim is to show that for every clause c in B , the gadget of exactly one of the three variables in c is in unstable state in M —this will translate to a 1-in-3 satisfying truth assignment for B .

Preferences will be set such that unstable (similarly, stable) states of gadgets in one level force a certain number of gadgets in the adjacent level to be in unstable (resp., stable) state. The reduction in [11] is also based on the same idea and for each clause in B , it used three level 0 gadgets, three level 2 gadgets, and one level 3 gadget (recall that clause gadgets are in levels 0, 2, 3). In our setting of complete preference lists, we will need “duplicates” or counterparts of all these gadgets to show the above reduction. So for each clause, we will use six level 0 gadgets, six level 2 gadgets, and two level 3 gadgets. In order to use Theorem 2, we will also need a new gadget to “glue” alternating paths across gadgets: this role will be performed by the Z -gadget which will be in stable state in M .

We will show that every level 3 gadget has to be in unstable state in M . Our technical lemma (Lemma 8) proves that this forces either two of the first three level 2 gadgets or two of the last three level 2 gadgets of every clause to be unstable state in M . We then show this forces at least one gadget of the three variables in every clause in B to be in unstable state in M .

We also show that every level 0 gadget has to be in stable state in M . This will induce at most one gadget of the three variables in every clause in B to be in unstable state in M . Thus *exactly one* gadget of the three variables in every clause in B will be in unstable state in M .

Our gadgets We will now describe all the gadgets that we use here: along with a figure, we provide the preference lists of vertices in this gadget. The tail of each list consists of all vertices not listed yet, in an arbitrary order. Even though the preference lists are complete, the structure of the gadgets and the preference lists will ensure that inter-gadget edges will not belong to any popular matching, as we will show in Section 4.

The D -gadget The D -gadget is on 4 vertices d_0, d_1, d_2, d_3 and the preference lists of these vertices are as given in Fig. 1 with all vertices outside the D -gadget at the tail of each list (in an arbitrary order). Recall that this gadget admits no stable matching. The role of D -gadget will be that of a *delimiter*—we will show that every vertex will have to be matched in any popular matching to a neighbor preferred to all its neighbors in the D -gadget.



Fig. 2 The variable gadget in level 1

We describe gadgets from level 1 first, then levels 0, 2, 3, and finally, the Z-gadget. The stable matchings within the gadgets are highlighted by colors in the figures. The gray elements in the preference lists denote vertices that are outside this gadget. We will assume that D in a preference list stands for $d_0 > d_1 > d_2 > d_3$.

Level 1 For each variable X_i in the formula B , we construct a gadget on four vertices as shown in Fig. 2.

The bottom vertices x'_i and y'_i will be preferred by some vertices in level 0 to vertices in their own gadget, while the top vertices x_i and y_i will be preferred by some vertices in level 2 to vertices in their own gadget. All four vertices in a level 1 gadget prefer to be matched among themselves, along the four edges drawn than be matched to any other vertex in the graph. This gadget has a unique stable matching $\{(x_i, y_i), (x'_i, y'_i)\}$.

Level 0 For each clause $c = X_i \vee X_j \vee X_k$ in the formula B , we create 6 gadgets in level 0. For every ordered pair of elements in $\{i, j, k\}$, there is one such gadget. One of these gadgets (this corresponds to the pair (j, k)) can be seen in Fig. 3. The top two vertices, i.e. a_1^c and b_1^c , rank y'_j and x'_k in level 1, as their respective second choices. Recall that indices j and k are well-defined in the clause $c = X_i \vee X_j \vee X_k$. Within this level 0 gadget on $a_1^c, b_1^c, a_2^c, b_2^c$, both $\{(a_1^c, b_1^c), (a_2^c, b_2^c)\}$ and $\{(a_1^c, b_2^c), (a_2^c, b_1^c)\}$ are stable matchings. In the preference lists below (and also for gadgets in levels 2 and 3), we have omitted the superscript c in their lists for the sake of readability.

The gadget on vertices $\{a_3^c, a_4^c, b_3^c, b_4^c\}$ is built analogously: the vertex a_3^c ranks y'_k as its second choice, while b_3^c ranks x'_i second. In the third gadget, the vertex a_5^c ranks y'_i second, while b_5^c ranks x'_j second. Observe the shift in i, j, k indices as second choices for vertices a_1^c, a_3^c, a_5^c (and similarly, for b_1^c, b_3^c, b_5^c).

The fourth, fifth, and sixth gadgets are analogous to their counterparts, the first, second, and third gadgets, respectively, but there is a slight twist. More precisely, the preferences of $a_1^{c'}, a_2^{c'}, b_1^{c'}, b_2^{c'}$ in the fourth gadget are analogous to the preferences in Fig. 3, except that $a_1^{c'}$ ranks y'_k second, while $b_1^{c'}$ ranks x'_j second.




Fig. 3 A clause gadget in level 0. The set of preference lists on the left belongs to the first gadget, while the set of preference lists on the right belongs to the fourth gadget (this corresponds to the pair (k, j))

Similarly, the second choice of a_3^c is y'_i , the second choice of b_3^c is x'_k , and finally, a_5^c ranks y'_j second, while b_5^c ranks x'_i second. Observe the change in *orientation* of the indices i, j, k as second choice neighbors when comparing the first three level 0 gadgets of c with its last three level 0 gadgets. This will be important to us later.

Level 2 For each clause $c = X_i \vee X_j \vee X_k$ in the formula B , we create 6 gadgets in level 2. The first gadget in level 2 is on vertices $p_0^c, p_1^c, p_2^c, q_0^c, q_1^c, q_2^c$ and their preference lists are described in Fig. 4. Note that p_2^c ranks y_j from level 1 as its second choice, while q_2^c ranks x_k from level 1 second.

The second gadget in level 2 is on vertices $p_3^c, p_4^c, p_5^c, q_3^c, q_4^c, q_5^c$ and it is built analogously. That is, p_3^c and q_3^c are each other's top choices and similarly, p_4^c and q_4^c are each other's top choices, and so on. The preference list of p_5^c is $q_3^c > y_k > q_4^c > q_5^c > D > \dots$ and the preference list of q_5^c is $p_4^c > x_i > p_3^c > p_5^c > D > \dots$

The third gadget in level 2 is on vertices $p_6^c, p_7^c, p_8^c, q_6^c, q_7^c, q_8^c$ and it is built analogously. In particular, the preference list of p_8^c is $q_6^c > y_i > q_7^c > q_8^c > D > \dots$ and the preference list of q_8^c is $p_7^c > x_j > p_6^c > p_8^c > D > \dots$

The fourth gadget in level 2 is on vertices $p_0^c, p_1^c, p_2^c, q_0^c, q_1^c, q_2^c$ and it is totally analogous to its counterpart, the first gadget in level 2. That is, p_0^c and q_0^c are each other's top choices and similarly, p_1^c and q_1^c are each other's top choices, and so on. In particular, the preference list of p_2^c is $q_0^c > y_j > q_1^c > q_2^c > D > \dots$ and the preference list of q_2^c is $p_1^c > x_k > p_0^c > p_2^c > D > \dots$

Similarly, the fifth gadget in level 2 is on vertices $p_3^c, p_4^c, p_5^c, q_3^c, q_4^c, q_5^c$ and it is totally analogous to the second gadget in level 2. Also, the sixth gadget in level 2 is on vertices $p_6^c, p_7^c, p_8^c, q_6^c, q_7^c, q_8^c$ and it is totally analogous to the third gadget in level 2.

Level 3 For each clause $c = X_i \vee X_j \vee X_k$ in the formula B , we create 2 gadgets in level 3. The first gadget is on vertices $s_0^c, s_1^c, s_2^c, s_3^c, t_0^c, t_1^c, t_2^c, t_3^c$ and the preference lists of these vertices are described in Fig. 5.

The counterpart of the first gadget in level 3 is the second gadget in level 3. It is on vertices $s_0^c, s_1^c, s_2^c, s_3^c, t_0^c, t_1^c, t_2^c, t_3^c$ and their preference lists are totally analogous to the preference lists of the first gadget in level 3.

The Z-gadget. The Z-gadget is on 6 vertices $z_0, z_1, z_2, z_3, z_4, z_5$ and the preference lists of these vertices are given in Fig. 6. The vertices in a set stand for all




Fig. 4 A clause gadget in level 2

$s_0 : t_1 > q_0 > t_2 > q_3 > t_3 > D > t_0 > \dots$
 $t_0 : s_3 > p_7 > s_2 > p_4 > s_1 > D > s_0 > \dots$
 $s_1 : t_1 > t_0 > D > \dots$
 $t_1 : s_1 > s_0 > D > \dots$
 $s_2 : t_2 > t_0 > D > \dots$
 $t_2 : s_2 > s_0 > D > \dots$
 $s_3 : t_3 > t_0 > D > \dots$
 $t_3 : s_3 > s_0 > D > \dots$




Fig. 5 A clause gadget in level 3

$z_0 : z_4 > z_5 > \cup_{i=1}^{\kappa} \{x_i, y_i\} > \cup_{i=0}^2 \cup_c \{p_{3i+1}^c, q_{3i}^c, p_{3i+1}^c, q_{3i}^c\} >$
 $\cup_{i=1}^6 \cup_c \{a_i^c, b_i^c, a_i^c, b_i^c\} > z_1 > z_2 > z_3 > D > \dots$
 $z_1 : z_5 > z_4 > \cup_{i=1}^{\kappa} \{x_i, y_i\} > \cup_{i=0}^2 \cup_c \{p_{3i+1}^c, q_{3i}^c, p_{3i+1}^c, q_{3i}^c\} >$
 $\cup_{i=1}^6 \cup_c \{a_i^c, b_i^c, a_i^c, b_i^c\} > z_0 > z_3 > z_2 > D > \dots$
 $z_2 : z_0 > z_1 > z_3 > z_4 > z_5 > D > \dots$
 $z_3 : z_1 > z_0 > z_2 > z_5 > z_4 > D > \dots$
 $z_4 : z_2 > z_3 > z_5 > z_0 > z_1 > D > \dots$
 $z_5 : z_3 > z_2 > z_4 > z_1 > z_0 > D > \dots$




Fig. 6 The Z-gadget

these vertices in a fixed arbitrary order. For example, $\cup_{i=1}^{\kappa} \{x_i, y_i\}$ denotes all the “top” vertices belonging to all the κ variable gadgets in a fixed arbitrary order.

Note that G is a complete graph on an even number of vertices and so every popular matching in G has to be a perfect matching.

4 Popular edges in G

Call an edge e in G popular if there is a popular matching M in G such that $e \in M$. In this section we identify edges that cannot be popular and show that every popular edge has to be an *intra-gadget* edge, i.e., it connects two vertices of the same gadget.

The following observation, which is straightforward, will be used repeatedly in our proofs.

Observation 2 *Let v be u 's top choice neighbor. If v is matched in M to a neighbor worse than u then (u, v) is a blocking edge to M .*

We now start restricting the set of edges that can possibly occur in a popular matching. Our first lemma eliminates some of the inter-gadget edges incident to vertices $s_0^c, t_0^c, s_0^c,$ and t_0^c in level 3.

Lemma 1 *For any clause c , no popular matching in G can match s_0^c (similarly, t_0^c) to a neighbor worse than t_0^c (resp., s_0^c). An analogous statement holds for $s_0^{t^c}$ and $t_0^{t^c}$.*

Proof Let M be a popular matching such that $(s_0^c, v) \in M$ for some vertex v such that $t_0^c > v$ in s_0^c 's list, i.e., s_0^c prefers t_0^c to v . We claim this implies:

- a (+, +) edge reachable from v via an alternating path in G_M that begins with a non-matching edge incident to v and
- a (+, +) edge reachable from s_0^c via an alternating path in G_M that begins with a non-matching edge incident to s_0^c .

If this is the same (+, +) edge then we have an alternating cycle in G_M with a (+, +) edge, a contradiction to M 's popularity (by Theorem 2). If these are two different (+, +) edges then there is an alternating path in G_M with two (+, +) edges, again a contradiction to M 's popularity (by Theorem 2).

- (1) If v is a top choice neighbor for some vertex (such as x_i, y_i, z_j for $1 \leq i \leq \kappa$, $0 \leq j \leq 5$ or d_1, d_2, d_3 or $a_\ell^r, b_\ell^r, p_{3h}^r, p_{3h+1}^r, q_{3h}^r, q_{3h+1}^r$ or their counterparts for $1 \leq \ell \leq 6$, $0 \leq h \leq 2$ and any clause r) then there is a (+, +) edge incident to v (by Observation 2).
- (2) Suppose v is one of $s_0^r, t_0^r, s_0^{r'}, t_0^{r'}$ for some clause r (note that $r \neq c$ in the case of s_0^r, t_0^r). Assume without loss of generality that $v = s_0^r$. Then either (s_0^r, t_0^r) is a (+, +) edge or t_0^r is matched in M to a neighbor better than s_0^r . Recall t_0^r 's preference list: every vertex that t_0^r prefers to s_0^r is either a top choice neighbor or it is d_0 . In the former case, there is a (+, +) edge incident to t_0^r 's partner (by Observation 2) and in the latter case also there is a (+, +) edge incident to d_0 since one of d_1, d_2, d_3 is matched to a neighbor worse than d_0 and so there is a (+, +) edge between this vertex in the set $\{d_1, d_2, d_3\}$ and d_0 . Since the edge (s_0^r, t_0^r) is a (+, -) edge, there is a (+, +) edge reachable from $v = s_0^r$ via an alternating path of length 2.
- (3) The only case left is when v is neither a top choice neighbor of some vertex nor one of $s_0^r, t_0^r, s_0^{r'}, t_0^{r'}$ for some clause r . So v is a vertex such as d_0 or x'_i, y'_i (for $1 \leq i \leq \kappa$) or $p_{3h+2}^r, q_{3h+2}^r, p_{3h+2}^{r'}, q_{3h+2}^{r'}$ (for $h = 0, 1, 2$ and some clause r). It is easy to see that there is a (+, +) edge reachable from v via an alternating path of length at most 2. For instance, either (x'_i, y'_i) is a (+, +) edge or $(x_i, y_i) \in M$ which creates the alternating path $(s_0^c, x'_i) - (y'_i, x_i) - (y_i, *)$, where (x_i, y_i) is a (+, +) edge.

Similarly, we can argue that there is a (+, +) edge reachable from s_0^c via an alternating path in G_M . If t_0^c is matched to a neighbor worse than s_0^c then the edge (s_0^c, t_0^c) is a (+, +) edge. Else t_0^c is matched to a neighbor u better than s_0^c and this means there is a (+, +) edge incident to u , as we argued above in case (2). Hence there is a (+, +) edge reachable from s_0^c via an alternating path of length at most 2 in G_M . □

Next we show that no inter-gadget edge incident to any vertex in the D -gadget can appear in any popular matching.

Lemma 2 Every popular matching matches the vertices in the D -gadget among themselves.

Proof Let M be a matching that matches d_i for some $i \in \{0, 1, 2, 3\}$ to a vertex v outside the D -gadget. This means at least 2 vertices d_i and d_j in the D -gadget are matched to vertices outside the D -gadget. So (d_i, d_j) is a $(+, +)$ edge. We now claim there is a forbidden alternating path or cycle (as given in Theorem 2) to M 's popularity.

If v is a top choice neighbor or a vertex such as x'_i, y'_i (for $1 \leq i \leq \kappa$) or $p_{3h+2}^c, q_{3h+2}^c, p'_{3h+2}, q'_{3h+2}$ (for $h = 0, 1, 2$ and some clause c) then there is a $(+, +)$ edge e reachable from v via an alternating path of length at most 2 as seen in the proof of Lemma 1. This creates an alternating path in G_M with 2 $(+, +)$ edges: (d_j, d_i) and e .

The other possibility is that v is $s_0^c, t_0^c, s_0^{t^c}, t_0^{s^c}$ for some clause c . Assume without loss of generality that $v = s_0^c$. Consider the vertex t_0^c . We know from Lemma 1 that t_0^c has to be matched to a neighbor at least as good as s_0^c . So we have the following cases:

- (1) t_0^c is matched to d_j in the D -gadget: this means there is either an alternating path with 2 $(+, +)$ edges or an alternating cycle with a $(+, +)$ edge:

$$(s_0^c, d_i) \overset{(+,+)}{\dashrightarrow} (d_j, t_0^c) \overset{(+,+)}{\dashrightarrow} (s_1^c, *) \quad \text{or}$$

$$(s_0^c, d_i) \overset{(+,+)}{\dashrightarrow} (d_j, t_0^c) \overset{(+,-)}{\dashrightarrow} (s_1^c, t_1^c) \overset{(-,+)}{\dashrightarrow} (s_0^c, d_i).$$

If s_1^c is matched to a neighbor worse than t_0^c then the former is an alternating path with two $(+, +)$ edges: these are (d_i, d_j) and (t_0^c, s_1^c) . Else s_1^c is matched to t_1^c and the latter is an alternating cycle with a $(+, +)$ edge, which is (d_i, d_j) .

- (2) t_0^c is matched to s_i^c for some $i \in \{1, 2, 3\}$: this means t_i^c is matched to a neighbor worse than s_0^c and so (s_0^c, t_i^c) is a $(+, +)$ edge and thus we have the following alternating path with two $(+, +)$ edges (t_i^c, s_0^c) and (d_i, d_j) :

$$(*, t_i^c) \overset{(+,+)}{\dashrightarrow} (s_0^c, d_i) \overset{(+,+)}{\dashrightarrow} (d_j, *).$$

- (3) t_0^c is matched to either p_4^c or p_7^c : we will show that this results in an alternating path with two $(+, +)$ edges. Assume without loss of generality that t_0^c is matched to p_4^c . Consider the following alternating path:

$$(*, s_3^c) \overset{(+,+)}{\dashrightarrow} (t_0^c, p_4^c) \overset{(+,+)}{\dashrightarrow} (q_5^c, *) \quad \text{or}$$

$$(*, d_j) \overset{(+,+)}{\dashrightarrow} (d_i, s_0^c) \overset{(+,-)}{\dashrightarrow} (t_3^c, s_3^c) \overset{(-,+)}{\dashrightarrow} (t_0^c, p_4^c) \overset{(+,+)}{\dashrightarrow} (q_5^c, *).$$

Recall that s_3^c is the top choice neighbor of t_0^c and the vertex p_4^c is the top choice neighbor of q_5^c . If the vertex s_3^c is matched to a neighbor worse than t_0^c then the former path is an alternating path in G_M with two $(+, +)$ edges in it: these are (s_3^c, t_0^c) and (p_4^c, q_5^c) . Else $(s_3^c, t_3^c) \in M$ and recall that the edge (s_0^c, t_3^c) is a $(+, -)$

edge as s_0^c prefers t_3^c to d_i . This creates the latter path which is an alternating path in G_M with 2 (+, +) edges in it: these are (d_i, d_j) and (p_4^c, q_5^c) . \square

The gadget D admits 2 popular matchings: $\{(d_0, d_1), (d_2, d_3)\}$ and $\{(d_0, d_2), (d_1, d_3)\}$. So if M is a popular matching then either $\{(d_0, d_1), (d_2, d_3)\} \subset M$ or $\{(d_0, d_2), (d_1, d_3)\} \subset M$. The following lemma further restricts the set of popular edges and this will be used repeatedly in our proof.

Lemma 3 *Let (u, v) be an edge in G where both u and v prefer d_0 to each other. Then (u, v) cannot be a popular edge.*

Proof Let M be a popular matching in G that contains such an edge (u, v) . We know from Lemma 2 that either $\{(d_0, d_1), (d_2, d_3)\} \subset M$ or $\{(d_0, d_2), (d_1, d_3)\} \subset M$. So there is always a blocking edge $(d_i, d_j) \in \{(d_1, d_3), (d_1, d_2)\}$ to M .

Observe that both u and v cannot belong to the D -gadget as there is no such pair within D . If exactly one of u, v belongs to the D -gadget then (u, v) is not a popular edge (by Lemma 2). So neither u nor v belongs to the D -gadget and this implies that u prefers d_0, d_1, d_2, d_3 to v and symmetrically, v prefers d_0, d_1, d_2, d_3 to u .

Consider the following alternating cycle C with respect to M :

$$(u, v) \overset{(+,-)}{\dashrightarrow} (d_i, d_i) \overset{(+,+)}{\dashrightarrow} (d_j, d_j) \overset{(-,+)}{\dashrightarrow} (u, v),$$

where (d_i, d_i) and (d_j, d_j) are edges from the D -gadget in M and (d_i, d_j) is a blocking edge. Thus C is an alternating cycle in G_M with a (+, +) edge. This contradicts the popularity of M (by Theorem 2). \square

Corollary 1 *The edges (s_0^c, t_0^c) and (s_0^c, t_0^c) are not popular edges for any clause c .*

Corollary 1 follows from Lemma 3 by setting u and v to s_0^c and t_0^c (similarly, s_0^c and t_0^c), respectively. Let us call u a level i vertex if u belongs to a level i gadget. Lemma 4 further restricts the set of popular edges; the proof of this lemma consists of three main claims.

Lemma 4 *No edge between a level i vertex and a level $i + 1$ vertex is popular, for $0 \leq i \leq 2$.*

The proof of Lemma 4 follows from Claims 1–3 proved below.

Claim 1 *There is no popular edge between a level 0 vertex and a level 1 vertex.*

Proof Let M be a popular matching in G with such an edge, say (a_1^c, y_j^c) , where $c = X_i \vee X_j \vee X_k$. We claim this would create an alternating path in G_M with two (+, +) edges in it (this would contradict Theorem 2). Consider the vertex b_1^c . There are 3 possibilities for b_1^c 's partner in M .

- (1) b_1^c is matched to a_2^c
 So (a_2^c, b_2^c) is labeled $(+, +)$. Recall that b_2^c is a_2^c 's top choice and the only neighbor that b_2^c prefers to a_2^c is a_1^c (which is matched to y_j'). Consider the following alternating path in G_M :

$$(*, b_2^c) \xrightarrow{(+,+)} (a_2^c, b_1^c) \xrightarrow{(-,+)} (a_1^c, y_j') \xrightarrow{(+,+)} (x_j', *).$$

If x_j' is matched to a neighbor worse than y_j' then the above is an alternating path in G_M with two $(+, +)$ edges: these are (b_2^c, a_2^c) and (y_j', x_j') . Else replace $(x_j', *)$ in the above path with $(x_j', y_j) - (x_j, *)$: the $(+, +)$ edges here are (b_2^c, a_2^c) and (y_j, x_j) .

- (2) b_1^c is matched to x_k'
 Either the edge (x_k', y_k') or the edge (x_k, y_k) will block M . Suppose y_k' is matched to a neighbor worse than x_k' in M . Consider the following alternating path in G_M :

$$(*, y_k') \xrightarrow{(+,+)} (x_k', b_1^c) \xrightarrow{(-,+)} (a_1^c, y_j') \xrightarrow{(+,+)} (x_j', *).$$

Either the above is an alternating path in G_M with two $(+, +)$ edges or by replacing $(x_j', *)$ with $(x_j', y_j) - (x_j, *)$ (as done in case (1)), we get an alternating path in G_M with two $(+, +)$ edges. If y_k' is matched to a neighbor better than x_k' in M , i.e., if $(x_k, y_k') \in M$ then prefix both these alternating paths with $(*, y_k)$. This will yield an alternating path in G_M with (x_k, y_k) as a blocking edge and either (x_j', y_j') or (x_j, y_j) as a blocking edge.

- (3) b_1^c is matched to a neighbor worse than a_1^c
 Then the edge (a_1^c, b_1^c) is labeled $(+, +)$. Consider the following alternating path in G_M :

$$\begin{aligned} (*, b_1^c) \xrightarrow{(+,+)} (a_1^c, y_j') \xrightarrow{(+,+)} (x_j', *) \quad \text{or} \\ (*, b_1^c) \xrightarrow{(+,+)} (a_1^c, y_j') \xrightarrow{(+,-)} (x_j', y_j) \xrightarrow{(+,+)} (x_j, *). \end{aligned}$$

That is, if x_j' is matched to a neighbor worse than y_j' then consider the first alternating path above: this is an alternating path in G_M with both (b_1^c, a_1^c) and (y_j', x_j') as $(+, +)$ edges. Else $(x_j', y_j) \in M$ and the second alternating path is an alternating path in G_M with (b_1^c, a_1^c) and (y_j, x_j) as $(+, +)$ edges.

Thus (a_1^c, y_j') cannot belong to any popular matching in G . Similarly, (b_1^c, x_k') also cannot belong to any popular matching. Suppose $(b_1^c, x_k') \in M$. Consider the vertex a_2^c : there are 2 possibilities for a_2^c 's partner in M .

- (1) a_2^c is matched to b_2^c
 Recall that a_1^c is b_2^c 's top choice and since a_1^c is not matched to either y_j' (by our proof above) or b_1^c (which is matched to x_k'), the edge (a_1^c, b_2^c) is labeled $(+, +)$. Consider the following alternating path in G_M :

$$(*, a_1^c) \overset{(+,+)}{\underset{-}} (b_2^c, a_2^c) \overset{(-,+)}{\underset{-}} (b_1^c, x'_k) \overset{(+,+)}{\underset{-}} (y'_k, *)$$

If y'_k is matched to a neighbor worse than x'_k then the above is an alternating path in G_M with two $(+, +)$ edges: these are (a_1^c, b_2^c) and (x'_k, y'_k) . Else replace $(y'_k, *)$ in the above path with $(y'_k, x_k) - (y_k, *)$: the $(+, +)$ edges here are (a_1^c, b_2^c) and (x_k, y_k) .

- (2) a_2^c is matched to a neighbor worse than b_1^c

Since a_2^c is b_1^c 's top choice, the edge (a_2^c, b_1^c) is labeled $(+, +)$. Consider the following alternating path in G_M :

$$(*, a_2^c) \overset{(+,+)}{\underset{-}} (b_1^c, x'_k) \overset{(+,+)}{\underset{-}} (y'_k, *) \quad \text{or} \\ (*, a_2^c) \overset{(+,+)}{\underset{-}} (b_1^c, x'_k) \overset{(+,-)}{\underset{-}} (y'_k, x_k) \overset{(+,+)}{\underset{-}} (y_k, *)$$

That is, if y'_k is matched to a neighbor worse than x'_k then consider the first alternating path above: this is an alternating path in G_M with both (a_2^c, b_1^c) and (x'_k, y'_k) as $(+, +)$ edges. Else $(y'_k, x_k) \in M$ and the second alternating path is an alternating path in G_M with (a_2^c, b_1^c) and (x_k, y_k) as $(+, +)$ edges.

Thus (b_1^c, x'_k) is also not a popular edge. The proof that $(a_3^c, y'_k), (a_5^c, y'_i), (a_1^c, y'_k), (a_3^c, y'_i), (a_5^c, y'_j)$ are not popular edges is analogous to the proof that (a_1^c, y'_j) is not a popular edge. Similarly, the proof that $(b_3^c, x'_i), (b_5^c, x'_j), (b_1^c, x'_j), (b_3^c, x'_k), (b_5^c, x'_i)$ are not popular edges is analogous to the proof that (b_1^c, x'_k) is not a popular edge. The presence of any other edge between a level 0 vertex corresponding to clause c and a level 1 vertex in the popular matching M would contradict Lemma 3. □

Claim 2 *There is no popular edge between a level 1 vertex and a level 2 vertex.*

Proof Let M be a popular matching in G that contains such an edge, say (p_2^c, y_j) , where $c = X_i \vee X_j \vee X_k$. Consider the following alternating path with respect to M :

$$(p_2^c, y_j) \overset{(+,+)}{\underset{-}} (x'_j, y'_j) \overset{(+,+)}{\underset{-}} (x_j, *)$$

Since M is a perfect matching, x'_j is matched in M . We know that no edge between x'_j and a level 0 vertex belongs to M (by Claim 1). Also, M cannot match x'_j to a neighbor that it regards worse than d_0 (by Lemma 3) or to any neighbor in the D -gadget (by Lemma 2). Thus x'_j has to be matched to y'_j in M and so the above alternating path has two $(+, +)$ edges: (x'_j, y_j) and (x_j, y'_j) . This is a contradiction to M 's popularity (by Theorem 2).

It can similarly be shown that (q_2^c, x_k) cannot belong to M . We consider the alternating path $(q_2^c, x_k) - (y'_k, x'_k) - (y_k, *)$ here to get a contradiction to Theorem 2.

The proof that M cannot include any other edge between a level 1 vertex and a level 2 vertex corresponding to clause c is either analogous to the above proof or the presence of such an edge in M violates Lemma 3. □

Claim 3 *There is no popular edge between a level 2 vertex and a level 3 vertex.*

Proof Let M be a popular matching in G that contains such an edge, say (s_0^c, q_0^c) , for some clause c . Consider the following alternating path with respect to M :

$$(s_0^c, q_0^c) \overset{(+,+)}{\dashrightarrow} (p_2^c, q_2^c) \overset{(+,+)}{\dashrightarrow} (p_0^c, *) \quad \text{or}$$

$$(s_0^c, q_0^c) \overset{(+,+)}{\dashrightarrow} (p_2^c, q_1^c) \overset{(+,+)}{\dashrightarrow} (p_1^c, *).$$

The vertex p_2^c is matched in M and its partner cannot be a level 1 vertex (by Claim 2) or a neighbor worse than d_0 (by Lemma 3) or to any neighbor in the D -gadget (by Lemma 2). So either (p_2^c, q_2^c) or (p_2^c, q_1^c) is in M . This means either the first alternating path given above or the second one is an alternating path in G_M with two $(+, +)$ edges: (p_2^c, q_0^c) and (p_0^c, q_2^c) in the former and (p_2^c, q_0^c) and (p_1^c, q_1^c) in the latter. This is a contradiction to M 's popularity (by Theorem 2).

The proof that M cannot include any other edge between a level 2 vertex and a level 3 vertex is either analogous to the above proof or the presence of such an edge in M violates Lemma 3. □

We now show that no popular matching contains an inter-gadget edge incident to any vertex in the Z -gadget. This is analogous to Lemma 2, which showed the same property for the D -gadget.

Lemma 5 *All popular matchings match the 6 vertices of the Z -gadget among themselves.*

Proof Let M be any popular matching in G . It follows from Lemma 3 that M has to match z_2, z_3, z_4 , and z_5 within the Z -gadget. Let us now show that z_0 also has to be matched within the Z -gadget. Then it immediately follows that z_1 also has to be matched within the Z -gadget. We have the following 3 cases:

- (1) Suppose z_0 is matched in M to a level 0 neighbor, say b_1^c , for some clause c . Then (a_1^c, b_1^c) is a blocking edge to M . Lemmas 2, 3, and 4 ensure that a_1^c is either matched to z_1 or to b_2^c . We investigate these two cases below.
 - $(a_1^c, z_1) \in M$: Here both z_0 and z_1 are matched to vertices they prefer to all their neighbors inside the Z -gadget, except for z_4 and z_5 . We know that z_4 and z_5 must be matched inside the Z -gadget. There are 3 subcases and in each case there is an alternating cycle in G_M with a blocking edge (a_1^c, b_1^c) : a contradiction to M 's popularity (by Theorem 2).
 - $(z_4, z_2) \in M$: the alternating cycle $(b_1^c, z_0) \overset{(+,-)}{\dashrightarrow} (z_4, z_2) \overset{(+,-)}{\dashrightarrow} (z_1, a_1^c) \overset{(+,+)}{\dashrightarrow} (b_1^c, z_0)$ is
 - $(z_4, z_3) \in M$: the alternating cycle $(b_1^c, z_0) \overset{(+,-)}{\dashrightarrow} (z_4, z_3) \overset{(+,-)}{\dashrightarrow} (z_1, a_1^c) \overset{(+,+)}{\dashrightarrow} (b_1^c, z_0)$ is
 - $(z_4, z_5) \in M$: the alternating cycle $(b_1^c, z_0) \overset{(+,-)}{\dashrightarrow} (z_4, z_5) \overset{(-,+)}{\dashrightarrow} (z_1, a_1^c) \overset{(+,+)}{\dashrightarrow} (b_1^c, z_0)$ is
 - $(a_1^c, b_2^c) \in M$: Lemmas 2, 3, and 4 ensure that a_2^c is matched to z_1 (recall that M is perfect). This leads to the same 3 subcases as above, except that instead

- of the edge (z_1, a_1^c) , there is the path $(z_1, a_2^c) - (b_2^c, a_1^c)$ in G_M : here (a_2^c, b_2^c) is labeled $(+, -)$.
- (2) Suppose z_0 is matched in M to a level 1 neighbor, say y_i , for some $i \in \{1, \dots, \kappa\}$. This case is similar to the previous case. Here the edge (x_i, y_i) becomes the blocking edge to M . It follows from Lemmas 2, 3, and 4 that x_i is either matched to z_1 or to y'_i . The latter case leaves x'_i unmatched and the subcases that arise in the former case are analogous to the ones in case (1).
 - (3) Suppose z_0 is matched in M to a level 2 neighbor, say q_0^c , for some clause c . It follows from Lemmas 2, 3, and 4 that (p_0^c, q_2^c) , (p_2^c, q_1^c) , and (p_1^c, z_1) are in M . Consider the alternating path $(z_0, q_0^c) - (p_2^c, q_1^c) - (p_1^c, z_1)$: it has two blocking edges (p_2^c, q_0^c) and (p_1^c, q_1^c) . This is again a contradiction to M 's popularity.

Recall that Lemma 2 showed that all vertices of the D -gadget must be matched within the gadget. Thus z_0 cannot be matched to a vertex in the D -gadget. The case where z_0 is matched in M to a level 3 neighbor does not arise as such an edge would violate Lemma 3. This finishes our proof that any popular matching M matches the 6 vertices of the Z -gadget among themselves. □

Our next lemma shows that the Z -gadget has a unique popular matching. The fact that the edge (z_0, z_1) has to be in any popular matching M will be used repeatedly in Section 5.

Lemma 6 *The only popular matching inside the Z -gadget is $\{(z_0, z_1), (z_2, z_3), (z_4, z_5)\}$.*

Proof The matching $\{(z_0, z_1), (z_2, z_3), (z_4, z_5)\}$ is stable in the Z -gadget, thus this is a popular matching. Note that this gadget has no other stable matching.

Let M be any popular matching. We know from Lemma 5 that M matches the 6 vertices of the Z -gadget among themselves. Suppose M contains one or more of the edges (z_i, z_j) where $i = j \pmod 2$ (colored black in Fig. 6).

This means one of the edges $(z_0, z_2), (z_2, z_4), (z_0, z_4)$ is in M . Let $(z_0, z_2) \in M$. There are three candidate matchings that we need to check for popularity: note that none is popular (by Theorem 2).

- $\{(z_0, z_2)_{(+,+)}, (z_1, z_3)_{(-,+)}\}$: this has the alternating cycle $(z_2, z_0)_{(+,-)} - (z_1, z_3)_{(-,+)} - (z_5, z_4)_{(+,-)} - (z_2, z_0)$ with the blocking edge (z_0, z_1) .
- $\{(z_0, z_2)_{(-,+)}, (z_1, z_4)_{(+,+)}\}$: this has the alternating cycle $(z_0, z_2)_{(-,+)} - (z_3, z_5)_{(-,+)} - (z_1, z_4)_{(+,+)} - (z_0, z_2)$ with the blocking edge (z_0, z_4) .
- $\{(z_0, z_2)_{(+,-)}, (z_1, z_5)_{(+,+)}\}$: this has the alternating cycle $(z_2, z_0)_{(+,-)} - (z_1, z_5)_{(+,+)} - (z_3, z_4)_{(+,-)} - (z_2, z_0)$ with the blocking edge (z_3, z_5) .

So (z_0, z_2) does not belong to M . We now claim that neither (z_2, z_4) nor (z_4, z_0) also belongs to M . This is because when confined to edges within the Z -gadget, the Z -gadget is symmetric with respect to z_0, z_2, z_4 (similarly, wrt z_1, z_3, z_5). Thus the same analysis as shown above for (z_0, z_2) holds for (z_2, z_4) and (z_4, z_0) as well by replacing every subscript s with $(s + 2) \pmod 6$ for the former edge and with $(s + 4) \pmod 6$ for the latter edge. Hence we can conclude that $M \subset \{(z_0, z_2, z_4)\} \times \{(z_1, z_3, z_5)\}$.

Suppose M contains an unstable edge here (dotted and gray in Fig. 6), say (z_0, z_3) : among the vertices in the Z -gadget, z_3 is the last choice of z_0 and the edge (z_0, z_2) blocks M . Since z_2 has to be matched in M , there are two cases.

- $(z_1, z_2) \in M$: the 4 vertices z_0, z_1, z_2, z_3 prefer $\{(z_0, z_2), (z_1, z_3)\}$ to $\{(z_0, z_3), (z_1, z_2)\} \subset M$.
- $(z_2, z_5) \in M$: the 4 vertices z_0, z_2, z_3, z_5 prefer $\{(z_0, z_2), (z_3, z_5)\}$ to $\{(z_0, z_3), (z_2, z_5)\} \subset M$.

Thus in both cases we have a contradiction to M 's popularity. Analogous proofs hold for other unstable edges chosen from $\{z_0, z_2, z_4\} \times \{z_1, z_3, z_5\}$. Thus the only popular matching inside the Z -gadget is $\{(z_0, z_1), (z_2, z_3), (z_4, z_5)\}$. \square

5 Stable states versus unstable states

In this section we will show how to obtain a 1-in-3 satisfying truth assignment for the input B from any popular matching in G . The following definition will be useful to us.

Definition 2 A gadget A in $G = (V, E)$ is said to be in *unstable state* with respect to matching M if there is a blocking edge $(u, v) \in V(A) \times V(A)$ with respect to M . If there is no such blocking edge to M then we say A is in *stable state* with respect to M .

In Figures 2, 3, 4, 5 and 6 depicting our gadgets, corresponding to matchings that consist of colored edges within the gadget, the relevant gadget is in *stable state*. A level 1 gadget in unstable state will encode the corresponding variable being set to true while a level 1 gadget in stable state will encode the corresponding variable being set to false. We will now analyze what gadgets are in stable/unstable state with respect to any popular matching M in G . This will lead to the proof that for any clause c , exactly one of the level 1 gadgets corresponding to the three variables in c is in unstable state.

Lemma 7 takes the starting step in this proof. Here M is any popular matching in G .

Lemma 7 For any clause c , the following statements hold:

- all its 6 level 0 gadgets are in stable state with respect to M ;
- both its level 3 gadgets in G are in unstable state with respect to M .

Proof Consider a level 0 gadget corresponding to clause c , say the one on vertices $a_1^c, b_1^c, a_2^c, b_2^c$. Lemmas 2, 3, 4, and 5 imply that either $\{(a_1^c, b_1^c), (a_2^c, b_2^c)\} \subset M$ or $\{(a_1^c, b_2^c), (a_2^c, b_1^c)\} \subset M$. Thus there is no blocking edge within this gadget. As this

holds for every level 0 gadget corresponding to c and for every clause c , the first part of the lemma follows.

We will now prove the second part of the lemma. Since M is a perfect matching, the vertices s_0^c, t_0^c (also $s_0^{t^c}, t_0^{t^c}$) have to be matched in M , for all clauses c . It follows from Lemmas 2 and 3 that both s_0^c and t_0^c (similarly, $s_0^{t^c}$ and $t_0^{t^c}$) have to be matched to neighbors that are better than d_0 . Lemma 4 showed that there is no popular edge between a level 3 vertex and a level 2 vertex. Thus s_0^c is matched to t_i^c for some $i \in \{1, 2, 3\}$.

If s_0^c is matched to t_i^c then s_i^c has to be matched to t_0^c —otherwise Lemma 3 would be violated by s_i^c and its partner. So (s_i^c, t_i^c) blocks M and this holds for every clause c . Similarly, there is a blocking edge (s_j^c, t_j^c) for some $j \in \{1, 2, 3\}$ for every clause c . □

Lemma 8 is our main technical lemma. This will be used in Lemma 9 to show that at least one of the level 1 gadgets corresponding to the three variables in clause c is in unstable state.

Lemma 8 *For any clause c , at least one of the following two conditions has to hold:*

- two or more of its first three level 2 gadgets are in unstable state with respect to M ;
- two or more of its last three level 2 gadgets are in unstable state with respect to M .

Proof Suppose both statements are false. Let M be a popular matching and c be a clause such that corresponding to clause c , at least two among its first three level 2 gadgets are in stable state with respect to M and at least two among its last three level 2 gadgets are in stable state with respect to M .

Consider the two level 3 gadgets corresponding to c . We know that $(s_0^c, t_i^c), (s_i^c, t_0^c)$ are in M for some $i \in \{1, 2, 3\}$ and similarly, $(s_0^{t^c}, t_j^{t^c}), (s_j^{t^c}, t_0^{t^c})$ are in M for some $j \in \{1, 2, 3\}$ (see the proof of Lemma 7). We will now show the existence of an alternating path ρ that will contradict M 's popularity.

For this, we claim it suffices to show that one gadget in each of the following two sets of gadgets is in *stable state*:

- the first three level 2 gadgets with a vertex that either s_0^c or t_0^c prefers to its partner in M ;
- the last three level 2 gadgets with a vertex that either $s_0^{t^c}$ or $t_0^{t^c}$ prefers to its partner in M .

For instance, suppose $i = 1$ and $j = 2$. So t_0^c prefers p_4^c and p_7^c to its partner s_1^c in M and s_0^c prefers $q_0^{t^c}$ to its partner $t_2^{t^c}$ in M and $t_0^{t^c}$ prefers $p_7^{t^c}$ to its partner $s_2^{t^c}$ in M . Consider the level 2 gadgets containing $p_4^c, p_7^c, q_0^{t^c}$, and $p_7^{t^c}$.

Observe that by our assumption in the first paragraph above, either the gadget of p_4^c or the gadget of p_7^c is in stable state, similarly either the gadget of q_0^c or the gadget of p_7^c is in stable state. In all 4 cases, we will show the existence of an alternating path ρ in G_M with two blocking edges (s_1^c, t_1^c) and (s_2^c, t_2^c) , which is a contradiction to M 's popularity (by Theorem 2).

1. Suppose the gadgets of p_4^c and q_0^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 3, 4, 5$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 0, 1, 2$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned} &(s_0^c, t_1^c) \overset{(+,+)}{\dashrightarrow} (s_1^c, t_0^c) \overset{(+,-)}{\dashrightarrow} (p_4^c, q_4^c) \overset{(-,+)}{\dashrightarrow} (p_5^c, q_5^c) \overset{(+,-)}{\dashrightarrow} (p_3^c, q_3^c) \overset{(-,+)}{\dashrightarrow} \\ &(z_0, z_1) \overset{(+,-)}{\dashrightarrow} (p_1^c, q_1^c) \overset{(-,+)}{\dashrightarrow} (p_2^c, q_2^c) \\ &\overset{(+,-)}{\dashrightarrow} (p_0^c, q_0^c) \overset{(-,+)}{\dashrightarrow} (s_0^c, t_2^c) \overset{(+,+)}{\dashrightarrow} (s_2^c, t_0^c). \end{aligned}$$

We know that $(z_0, z_1) \in M$ (by Lemma 6). Note that ρ is the desired alternating path in G_M with two blocking edges (s_1^c, t_1^c) and (s_2^c, t_2^c) .

2. Suppose the gadgets of p_4^c and p_7^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 3, 4, 5$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 6, 7, 8$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned} &(s_0^c, t_1^c) \overset{(+,+)}{\dashrightarrow} (s_1^c, t_0^c) \overset{(+,-)}{\dashrightarrow} (p_4^c, q_4^c) \overset{(-,+)}{\dashrightarrow} (p_5^c, q_5^c) \overset{(+,-)}{\dashrightarrow} (p_3^c, q_3^c) \overset{(-,+)}{\dashrightarrow} \\ &(z_0, z_1) \overset{(+,-)}{\dashrightarrow} (q_6^c, p_6^c) \overset{(-,+)}{\dashrightarrow} (q_8^c, p_8^c) \\ &\overset{(+,-)}{\dashrightarrow} (q_7^c, p_7^c) \overset{(-,+)}{\dashrightarrow} (t_0^c, s_2^c) \overset{(+,+)}{\dashrightarrow} (t_2^c, s_0^c). \end{aligned}$$

Observe that the labels on edges of $\rho \setminus M$ are identical to the first case and thus ρ is the desired alternating path in G_M with two blocking edges (s_1^c, t_1^c) and (s_2^c, t_2^c) .

3. Suppose the gadgets of p_7^c and q_0^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 6, 7, 8$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 0, 1, 2$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned} &(s_0^c, t_1^c) \overset{(+,+)}{\dashrightarrow} (s_1^c, t_0^c) \overset{(+,-)}{\dashrightarrow} (p_7^c, q_7^c) \overset{(-,+)}{\dashrightarrow} (p_8^c, q_8^c) \overset{(+,-)}{\dashrightarrow} (p_6^c, q_6^c) \overset{(-,+)}{\dashrightarrow} \\ &(z_0, z_1) \overset{(+,-)}{\dashrightarrow} (p_1^c, q_1^c) \overset{(-,+)}{\dashrightarrow} (p_2^c, q_2^c) \\ &\overset{(+,-)}{\dashrightarrow} (p_0^c, q_0^c) \overset{(-,+)}{\dashrightarrow} (s_0^c, t_2^c) \overset{(+,+)}{\dashrightarrow} (s_2^c, t_0^c). \end{aligned}$$

Again, observe that the labels on edges of $\rho \setminus M$ are identical to the first two cases and ρ is the desired alternating path with two blocking edges (s_1^c, t_1^c) and (s_2^c, t_2^c) .

4. Suppose the gadgets of p_7^c and p_7^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 6, 7, 8$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 6, 7, 8$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned}
 &(s_0^c, t_1^c)^{(+,+)} (s_1^c, t_0^c)^{(+,-)} (p_7^c, q_7^c)^{(-,+)} (p_8^c, q_8^c)^{(+,-)} (p_6^c, q_6^c)^{(-,+)} \\
 &(z_0, z_1)^{(+,-)} (q_6^c, p_6^c)^{(-,+)} (q_8^c, p_8^c) \\
 &^{(+,-)} (q_7^c, p_7^c)^{(-,+)} (t_0^c, s_2^c)^{(+,+)} (t_2^c, s_0^c).
 \end{aligned}$$

As before, the labels on edges of $\rho \setminus M$ are identical to the above three cases and ρ is the desired alternating path with two blocking edges (s_1^c, t_1^c) and (s_2^c, t_2^c) .

For any $(i, j) \in \{1, 2, 3\} \times \{1, 2, 3\}$, an analogous construction can be shown.

- Let $i = j = 1$. So t_0^c prefers p_4^c and p_7^c to its partner s_1^c in M and t_0^c prefers p_4^c and p_7^c to its partner s_1^c in M . We know that either the gadget of p_4^c or the gadget of p_7^c is in stable state, and similarly, either the gadget of p_4^c or the gadget of p_7^c is in stable state. Suppose the gadgets of p_4^c and p_7^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 3, 4, 5$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 3, 4, 5$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned}
 &(s_0^c, t_1^c)^{(+,+)} (s_1^c, t_0^c)^{(+,-)} (p_4^c, q_4^c)^{(-,+)} (p_5^c, q_5^c)^{(+,-)} (p_3^c, q_3^c)^{(-,+)} \\
 &(z_0, z_1)^{(+,-)} (q_3^c, p_3^c)^{(-,+)} (q_5^c, p_5^c) \\
 &^{(+,-)} (q_4^c, p_4^c)^{(-,+)} (t_0^c, s_1^c)^{(+,+)} (t_1^c, s_0^c).
 \end{aligned}$$

Observe again that the labels on edges of $\rho \setminus M$ are identical to the labels obtained for the desired alternating paths when $i = 1$ and $j = 2$. The path ρ is the desired alternating path in G_M with two blocking edges (s_1^c, t_1^c) and (s_1^c, t_1^c) . The case when the gadgets of p_7^c and p_7^c are in stable state was already seen in Case 4 of $i = 1$ and $j = 2$. The only difference between the path that we will construct now with the path ρ seen there is in the last two edges: now we will have (t_0^c, s_1^c) and (t_1^c, s_0^c) in M ; thus the blocking edges to our path will be (s_1^c, t_1^c) and (s_1^c, t_1^c) . The proofs for the remaining two cases: (i) when the gadgets of p_4^c and p_7^c are in stable state and (ii) when the gadgets of p_7^c and p_4^c are in stable state are totally analogous to the above case. Thus in all 4 cases, we can show the existence of an alternating path ρ in G_M with two blocking edges (s_1^c, t_1^c) and (s_1^c, t_1^c) : a contradiction to M 's popularity.

- Let $i = 1$ and $j = 3$. So t_0^c prefers p_4^c and p_7^c to its partner s_1^c in M and s_0^c prefers q_0^c and q_3^c to its partner t_3^c in M . We know that either the gadget of p_4^c or the gadget of p_7^c is in stable state, and similarly, either the gadget of q_0^c or the gadget of q_3^c is in stable state. The cases when the gadgets of p_4^c and q_0^c are in stable state and when the gadgets of p_7^c and q_0^c are in stable state were already seen in Case 1 and Case 3 of $i = 1$ and $j = 2$: thus we can construct analogous alternating paths in these cases. So let us consider the case when the gadgets of p_7^c and q_3^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 6, 7, 8$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 3, 4, 5$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned}
 & (s_0^c, t_1^c)^{(+,+)} (s_1^c, t_0^c)^{(+,-)} (p_7^c, q_7^c)^{(-,+)} (p_8^c, q_8^c)^{(+,-)} (p_6^c, q_6^c)^{(-,+)} \\
 & (z_0, z_1)^{(+,-)} (p_4^c, q_4^c)^{(-,+)} (p_5^c, q_5^c)^{(-,+)} \\
 & (p_3^c, q_3^c)^{(+,-)} (s_0^c, t_3^c)^{(-,+)} (s_3^c, t_0^c)^{(+,+)} .
 \end{aligned}$$

The above path ρ is the desired alternating path in G_M with two blocking edges (s_1^c, t_1^c) and (s_3^c, t_3^c) . The remaining case, i.e., when the gadgets of p_4^c and q_3^c are in stable state, is totally analogous to the above case and we can again show an alternating path in G_M with two blocking edges (s_1^c, t_1^c) and (s_3^c, t_3^c) .

- Let $i = 2$ and $j = 1$. This is a “mirror image” of the very first case considered: when $i = 1$ and $j = 2$. The only difference is that we swap *primed variables* and *unprimed variables* in ρ . For example, when the gadgets of q_0^c and p_4^c are in stable state, the desired alternating path is exactly the same as ρ in Case 1 there, except for this swapping of roles. Thus, ρ with blocking edges (s_2^c, t_2^c) and (s_1^c, t_1^c) , would be:

$$\begin{aligned}
 & (t_0^c, s_2^c)^{(+,+)} (t_2^c, s_0^c)^{(+,-)} (q_0^c, p_0^c)^{(-,+)} (q_2^c, p_2^c)^{(+,-)} (q_1^c, p_1^c)^{(-,+)} \\
 & (z_0, z_1)^{(+,-)} (q_3^c, p_3^c)^{(-,+)} (q_5^c, p_5^c)^{(-,+)} \\
 & (p_4^c, p_4^c)^{(+,-)} (t_0^c, s_1^c)^{(-,+)} (t_1^c, s_0^c)^{(+,+)} .
 \end{aligned}$$

- Let $i = j = 2$. So s_0^c prefers q_0^c to its partner t_2^c in M and t_0^c prefers p_7^c to its partner s_2^c in M and s_0^c prefers q_0^c to its partner t_2^c in M and t_0^c prefers p_7^c to its partner s_1^c in M . We know that either the gadget of q_0^c or the gadget of p_7^c is in stable state, and similarly, either the gadget of q_0^c or the gadget of p_7^c is in stable state. The cases when the gadgets of p_7^c and q_0^c are in stable state and when the gadgets of p_7^c and p_7^c are in stable state were already seen in Cases 3 and 4 of $i = 1$ and $j = 2$. Let us consider the case when the gadgets of q_0^c and q_0^c are in stable state. So the edges $(p_\ell^c, q_\ell^c) \in M$ for $\ell = 0, 1, 2$ and the edges $(p_h^c, q_h^c) \in M$ for $h = 0, 1, 2$. Consider the following alternating path ρ with respect to M :

$$\begin{aligned}
 & (t_0^c, s_2^c)^{(+,+)} (t_2^c, s_0^c)^{(+,-)} (q_0^c, p_0^c)^{(-,+)} (q_2^c, p_2^c)^{(+,-)} (q_1^c, p_1^c)^{(-,+)} \\
 & (z_0, z_1)^{(+,-)} (p_1^c, q_1^c)^{(-,+)} (p_2^c, q_2^c)^{(-,+)} \\
 & (p_0^c, q_0^c)^{(+,-)} (s_0^c, t_2^c)^{(-,+)} (s_2^c, t_0^c)^{(+,+)} .
 \end{aligned}$$

The path ρ is the desired alternating path in G_M with two blocking edges (s_2^c, t_2^c) and (s_2^c, t_2^c) . The proof for the remaining case is totally analogous. Thus in all 4 cases, we can show the existence of an alternating path ρ in G_M with two blocking edges (s_2^c, t_2^c) and (s_2^c, t_2^c) : a contradiction to M 's popularity. It is easy to see that the remaining cases of (i, j) are totally analogous to the ones listed above and this finishes the proof of the lemma. □

Recall that there are three level 1 gadgets associated with any clause c : these gadgets correspond to the three variables in c .

Lemma 9 *Let $c = X_i \vee X_j \vee X_k$. At least one of the level 1 gadgets corresponding to X_i, X_j, X_k is in unstable state with respect to M .*

Proof Suppose not. That is, assume that for some clause c , all three of its level 1 gadgets are in stable state. Let $c = X_i \vee X_j \vee X_k$. So (x_r, y_r) and (x'_r, y'_r) are in M for all $r \in \{i, j, k\}$.

We know from Lemma 8 that either two or more of the *first* three level 2 gadgets corresponding to c are in unstable state with respect to M ; or two or more of the *last* three level 2 gadgets corresponding to c are in unstable state with respect to M . We assume without loss of generality that the first and second gadgets, i.e., those on p_i^c, q_i^c , for $0 \leq i \leq 5$, are in unstable state with respect to M .

We know from our lemmas in Section 4 that there is no popular edge across gadgets. Thus M matches the 6 vertices of a level 2 gadget with each other. In particular, it follows from Lemma 3 that for the level 2 gadget on p_i^c, q_i^c for $i = 0, 1, 2$, we have (i) $(p_0^c, q_0^c), (p_1^c, q_1^c), (p_2^c, q_2^c)$ in M or (ii) $(p_0^c, q_2^c), (p_1^c, q_1^c), (p_2^c, q_0^c)$ in M or (iii) $(p_0^c, q_0^c), (p_1^c, q_2^c), (p_2^c, q_1^c)$ in M .

There are two unstable states for each level 2 gadget, i.e., either (ii) or (iii) above for the gadget on p_i^c, q_i^c for $i = 0, 1, 2$. A level 2 gadget can be in either of these two unstable states in M —without loss of generality assume that M contains $(p_0^c, q_0^c), (p_1^c, q_2^c), (p_2^c, q_1^c)$ and $(p_3^c, q_5^c), (p_4^c, q_4^c), (p_5^c, q_3^c)$. Observe that p_2^c likes y_j more than q_1^c and similarly, q_5^c likes x_i more than p_3^c . Consider the following alternating path ρ with respect to M :

$$(q_2^c, p_1^c) \overset{(+,+)}{\dashrightarrow} (q_1^c, p_2^c) \overset{(+,-)}{\dashrightarrow} (y_j, x_i) \overset{(-,+)}{\dashrightarrow} (z_0, z_1) \overset{(+,-)}{\dashrightarrow} (y_i, x_i) \overset{(-,+)}{\dashrightarrow} (q_5^c, p_3^c) \overset{(+,+)}{\dashrightarrow} (q_3^c, p_5^c).$$

Note that M has to contain (z_0, z_1) (by Lemma 6). Observe that ρ is an alternating path in G_M with *two* blocking edges (p_1^c, q_1^c) and (p_3^c, q_3^c) . This is a contradiction to M 's popularity (by Theorem 2) and the lemma follows. \square

Lemma 10 *Let $c = X_i \vee X_j \vee X_k$. At most one of the level 1 gadgets corresponding to X_i, X_j, X_k is in unstable state with respect to M .*

Proof Suppose not. So at least two of the three level 1 gadgets corresponding to X_i, X_j, X_k are in unstable state with respect to M . Assume without loss of generality that the gadgets corresponding to variables X_i and X_j are in unstable state. So the edges $(x_i, y'_i), (x'_i, y_i)$ are in M , similarly the edges $(x_j, y'_j), (x'_j, y_j)$ are in M .

Recall that a_5^c regards y'_i as its second choice neighbor and b_5^c regards x'_j as its second choice neighbor. Similarly, b_5^c regards x'_i as its second choice neighbor and a_5^c regards y'_j as its second choice neighbor.

In the popular matching M , level 0 vertices are matched within their own gadget. Therefore, either $\{(a_5^c, b_5^c), (a_6^c, b_6^c)\} \subset M$ or $\{(a_5^c, b_6^c), (a_6^c, b_5^c)\} \subset M$; similarly,

$\{(a_5^c, b_5^c), (a_6^c, b_6^c)\} \subset M$ or $\{(a_5^c, b_6^c), (a_6^c, b_5^c)\} \subset M$. Thus, the following two observations clearly hold:

- either a_5^c or b_5^c is matched to its third choice neighbor;
- either a_5^c or b_5^c is matched to its third choice neighbor.

Based on which of these vertices are matched to their third choice neighbors, we have four cases as shown below. Each of these 4 cases results in a forbidden alternating path/cycle (as given in Theorem 2), thus proving the lemma.

Case 1 The vertices a_5^c and a_5^c are matched to their third choice neighbors. So $(a_5^c, b_6^c), (a_6^c, b_5^c)$ and $(a_5^c, b_6^c), (a_6^c, b_5^c)$ are in M . Consider the following alternating path ρ with respect to M :

$$(x'_i, y_i) \overset{(+,+)}{\dashrightarrow} (x_i, y'_i) \overset{(-,+)}{\dashrightarrow} (a_5^c, b_6^c) \overset{(-,+)}{\dashrightarrow} (z_0, z_1) \overset{(+,-)}{\dashrightarrow} (b_6^c, a_5^c) \overset{(+,-)}{\dashrightarrow} (y'_j, x_j) \overset{(+,+)}{\dashrightarrow} (y_j, x'_j).$$

Observe that ρ is an alternating path in G_M with two blocking edges (x_i, y_i) and (x_j, y_j) , a contradiction to M 's popularity.

Case 2 The vertices a_5^c and b_5^c are matched to their third choice neighbors. So $(a_5^c, b_6^c), (a_6^c, b_5^c)$ and $(a_5^c, b_6^c), (a_6^c, b_5^c)$ are in M . Consider the following alternating cycle C with respect to M :

$$(b_6^c, a_5^c) \overset{(+,-)}{\dashrightarrow} (y'_i, x_i) \overset{(+,+)}{\dashrightarrow} (y_i, x'_i) \overset{(-,+)}{\dashrightarrow} (b_5^c, a_5^c) \overset{(-,+)}{\dashrightarrow} (z_1, z_0) \overset{(+,-)}{\dashrightarrow} (b_6^c, a_5^c).$$

Observe that C is an alternating cycle in G_M with a blocking edge (x_i, y_i) , a contradiction to M 's popularity.

Case 3 The vertices b_5^c and a_5^c are matched to their third choice neighbors. So $(a_5^c, b_5^c), (a_6^c, b_6^c)$ and $(a_5^c, b_6^c), (a_6^c, b_5^c)$ are in M . Consider the following alternating cycle C' with respect to M :

$$(a_5^c, b_5^c) \overset{(+,-)}{\dashrightarrow} (x'_j, y_j) \overset{(+,+)}{\dashrightarrow} (x_j, y'_j) \overset{(-,+)}{\dashrightarrow} (a_5^c, b_6^c) \overset{(-,+)}{\dashrightarrow} (z_0, z_1) \overset{(+,-)}{\dashrightarrow} (a_5^c, b_5^c).$$

Observe that C' is an alternating cycle in G_M with a blocking edge (x_j, y_j) , a contradiction to M 's popularity.

Case 4 The vertices b_5^c and b_5^c are matched to their third choice neighbors. So $(a_5^c, b_5^c), (a_6^c, b_6^c)$ and $(a_5^c, b_6^c), (a_6^c, b_5^c)$ are in M . Consider the following alternating path ρ' with respect to M :

$$(y'_i, x_i) \overset{(+,+)}{\dashrightarrow} (y_i, x'_i) \overset{(-,+)}{\dashrightarrow} (b_5^c, a_5^c) \overset{(-,+)}{\dashrightarrow} (z_0, z_1) \overset{(+,-)}{\dashrightarrow} (a_5^c, b_5^c) \overset{(+,-)}{\dashrightarrow} (x'_j, y_j) \overset{(+,+)}{\dashrightarrow} (x_j, y'_j).$$

Observe that ρ' is an alternating path in G_M with two blocking edges (x_i, y_i) and (x_j, y_j) , a contradiction to M 's popularity. □

Let M be a popular matching in G . It follows from Lemmas 9 and 10 that *exactly one* of the level 1 gadgets corresponding to the variables in c , for every clause c in B , is in unstable state with respect to M . This allows us to set a 1-in-3 satisfying truth assignment to instance B . For each variable X_i in B do:

- if the gadget corresponding to X_i is in *unstable* state then set $X_i = \text{true}$ else set $X_i = \text{false}$.

It follows from Lemmas 9 and 10 that this is a 1-in-3 satisfying truth assignment for B . We have thus shown the following result.

Theorem 3 *If G admits a popular matching then B has a 1-in-3 satisfying truth assignment.*

6 The converse

We will now show the converse of Theorem 3, i.e., if B has a 1-in-3 satisfying truth assignment S then G admits a popular matching. We will use S to construct a popular matching M in G as follows. To begin with, $M = \emptyset$.

Level 1 For each variable X_i do:

- if X_i is set to true in S then add (x_i, y'_i) and (x'_i, y_i) to M ;
- else add (x_i, y_i) and (x'_i, y'_i) to M .

Remark Note that the level 1 gadget of a variable set to true is in unstable state and the level 1 gadget of a variable set to false is in stable state.

For each clause $c = X_i \vee X_j \vee X_k$, we know that exactly one of X_i, X_j, X_k is set to true in S . Assume without loss of generality that $X_k = \text{true}$ in S . For the level 0, 2, and 3 gadgets corresponding to c , we do as follows:

Level 0 Recall that there are six level 0 gadgets that correspond to c . For the first 3 gadgets (these are on vertices a_i^c, b_i^c for $i = 1, \dots, 6$) do:

- include $(a_1^c, b_2^c), (a_2^c, b_1^c)$ from the first gadget;
- include $(a_3^c, b_3^c), (a_4^c, b_4^c)$ from the second gadget;
- choose either $(a_5^c, b_5^c), (a_6^c, b_6^c)$ or $(a_5^c, b_6^c), (a_6^c, b_5^c)$ from the third gadget.

Observe that since the third variable X_k of c was set to be true, cross edges are fixed in the first gadget (see Fig. 3), while the other stable matching (horizontal edges) is chosen in the second gadget.

For the fourth and fifth gadgets, we will do exactly the opposite. Also, it will not matter which stable pair of edges is chosen from the third and sixth gadgets.

So for the last 3 level 0 gadgets corresponding to c (these are on vertices $a_i^{l_c}, b_i^{l_c}$ for $i = 1, \dots, 6$) do:

- include $(a_1^{l_c}, b_1^{l_c}), (a_2^{l_c}, b_2^{l_c})$ from the fourth gadget;
- include $(a_3^{l_c}, b_4^{l_c}), (a_4^{l_c}, b_3^{l_c})$ from the fifth gadget.
- choose either $(a_5^{l_c}, b_5^{l_c}), (a_6^{l_c}, b_6^{l_c})$ or $(a_5^{l_c}, b_6^{l_c}), (a_6^{l_c}, b_5^{l_c})$ from the sixth gadget.

Level 2 Recall that there are six level 2 gadgets that correspond to c . For the first 3 gadgets (these are on vertices p_i^c, q_i^c for $i = 0, \dots, 8$) do:

- include $(p_0^c, q_2^c), (p_1^c, q_1^c), (p_2^c, q_0^c)$ from the first gadget
- include $(p_3^c, q_3^c), (p_4^c, q_5^c), (p_5^c, q_4^c)$ from the second gadget
- include $(p_6^c, q_6^c), (p_7^c, q_7^c), (p_8^c, q_8^c)$ from the third gadget

In the first three gadgets, because $X_k = \text{true}$, the third one is set to parallel edges, reaching the stable state, while the first one is blocked by the top horizontal edge and the second one is blocked by the middle horizontal edge. Include isomorphic edges (to the above ones) from the last three level 2 gadgets corresponding to c , i.e., include $(p_0^{l_c}, q_2^{l_c}), (p_1^{l_c}, q_1^{l_c}), (p_2^{l_c}, q_0^{l_c})$ from the fourth gadget, and so on. On this level, the last three gadgets mimic the matching edges from the first three gadgets, unlike in level 0.

Level 3 For the first level 3 gadget corresponding to c do:

- include $(s_0^c, t_3^c), (s_1^c, t_1^c), (s_2^c, t_2^c), (s_3^c, t_0^c)$ in M .

Since the third variable in c was set to be true, the vertices s_0^c and t_0^c are matched to t_3^c and s_3^c , respectively—thus the bottom horizontal edge (s_3^c, t_3^c) blocks M . Include isomorphic edges (to the above ones) for the second level 3 gadget corresponding to c , i.e., include $(s_0^{l_c}, t_3^{l_c}), (s_1^{l_c}, t_1^{l_c}), (s_2^{l_c}, t_2^{l_c}), (s_3^{l_c}, t_0^{l_c})$ in M . Once again, the second gadget mimics the matching edges on the first gadget.

Z-gadget and D-gadget. Finally include the edges $(z_0, z_1), (z_2, z_3), (z_4, z_5)$ from the Z -gadget in M . By Lemma 6, every popular matching in G has to include these edges. Also include the edges $(d_0, d_1), (d_2, d_3)$ from the D -gadget in M .

6.1 The popularity of M

We will now prove the popularity of the above matching M via the LP framework of popular matchings initiated in [22] for bipartite graphs. This framework generalizes to provide a sufficient condition for popularity in non-bipartite graphs [11]. This involves showing a witness $\alpha \in \mathbb{R}^n$ such that α is a *certificate* of M 's popularity. In order to define the constraints that α has to satisfy so as to certify M 's popularity, let us define an edge weight function w_M as follows.

For any edge (u, v) in G do:

- if (u, v) is labeled $(-, -)$ then set $w_M(u, v) = -2$;

- if (u, v) is labeled $(+, +)$ then set $w_M(u, v) = 2$;
- else set $w_M(u, v) = 0$. (So $w_M(e) = 0$ for all $e \in M$.)

Let N be any perfect matching in G . It is easy to see from the definition of the edge weight function w_M that $w_M(N) = \phi(N, M) - \phi(M, N)$.

Let the max-weight perfect fractional matching LP in the graph G with edge weight function w_M be our primal LP. This is **LP1** defined below. Here $\delta(u)$ denotes the set of edges incident to vertex u .

$$\begin{aligned}
 &\text{maximize } \sum_{e \in E} w_M(e)x_e \\
 &\text{subject to} \\
 &\sum_{e \in \delta(u)} x_e = 1 \quad \forall u \in V \quad \text{and} \quad x_e \geq 0 \quad \forall e \in E
 \end{aligned}
 \tag{LP1}$$

If the optimal value of **LP1** is at most 0 then $w_M(N) \leq 0$ for all perfect matchings N in G , i.e., $\phi(N, M) \leq \phi(M, N)$. Observe that this means $\phi(M', M) \leq \phi(M, M')$ for all matchings M' in G . This is because G is a complete graph on an even number of vertices, so for any matching M' , there is a perfect matching N such that $M' \subseteq N$. Thus $\phi(M', M) \leq \phi(N, M) \leq \phi(M, N) \leq \phi(M, M')$. Hence M is a popular matching in G .

Consider the LP that is dual to **LP1**. This is **LP2** given below in variables α_u , where $u \in V$.

$$\begin{aligned}
 &\text{minimize } \sum_{u \in V} \alpha_u \\
 &\text{subject to} \\
 &\alpha_u + \alpha_v \geq w_M(u, v) \quad \forall (u, v) \in E
 \end{aligned}
 \tag{LP2}$$

If we show a dual feasible solution α such that $\sum_{u \in V} \alpha_u = 0$ then the primal optimal value is at most 0, i.e., M is a popular matching. In order to prove the popularity of M , we define α as follows. For $r \in \{1, \dots, \kappa\}$ do: (recall that κ is the number of variables in the formula B)

- if X_r was set to true then set $\alpha_{x_r} = \alpha_{y_r} = 1$ and $\alpha_{x'_r} = \alpha_{y'_r} = -1$;
- else set $\alpha_{x_r} = \alpha_{y_r} = \alpha_{x'_r} = \alpha_{y'_r} = 0$.

Let clause $c = X_i \vee X_j \vee X_k$. Recall that we assumed that $X_i = X_j = \text{false}$ and $X_k = \text{true}$. For the vertices in clauses corresponding to c , we will set α -values as follows.

- For every level 0 vertex v do: set $\alpha_v = 0$.
- For the first three level 2 gadgets corresponding to c do:
 - set $\alpha_{p_0^c} = \alpha_{q_0^c} = 1$, $\alpha_{p_1^c} = 1$, $\alpha_{q_1^c} = -1$, and $\alpha_{p_2^c} = \alpha_{q_2^c} = -1$;
 - set $\alpha_{p_3^c} = -1$, $\alpha_{q_3^c} = 1$, $\alpha_{p_4^c} = \alpha_{q_4^c} = 1$, and $\alpha_{p_5^c} = \alpha_{q_5^c} = -1$;
 - set $\alpha_{p_6^c} = \alpha_{q_6^c} = \alpha_{p_7^c} = \alpha_{q_7^c} = \alpha_{p_8^c} = \alpha_{q_8^c} = 0$.

The setting of α -values is analogous for vertices in the last three level 2 gadgets corresponding to c . For the first level 3 gadget corresponding to c do:

- set $\alpha_{s_0^c} = \alpha_{t_0^c} = -1, \alpha_{s_1^c} = -1, \alpha_{t_1^c} = 1, \alpha_{s_2^c} = -1, \alpha_{t_2^c} = 1$, and $\alpha_{s_3^c} = \alpha_{t_3^c} = 1$.

The setting of α -values is analogous for vertices in the other level 3 gadget corresponding to c . For the z -vertices do: set $\alpha_u = 0$ for all $u \in \{z_0, \dots, z_5\}$. For the d -vertices do:

- set $\alpha_{d_0} = \alpha_{d_2} = -1$ and $\alpha_{d_1} = \alpha_{d_3} = 1$.

Properties of α . For every $(u, v) \in M$, either $\alpha_u = \alpha_v = 0$ or $\{\alpha_u, \alpha_v\} = \{-1, 1\}$; so $\alpha_u + \alpha_v = 0$. Since M is a perfect matching, we have $\sum_{u \in V} \alpha_u = 0$. The claims stated below show that α is a feasible solution to LP2. This will prove the popularity of M .

We need to show that every edge (u, v) is covered, i.e., $\alpha_u + \alpha_v \geq w_M(u, v)$. We have already observed that for any $(u, v) \in M, \alpha_u + \alpha_v = 0 = w_M(u, v)$.

Claim 4 *Let (u, v) be an intra-gadget blocking edge to M . Then $\alpha_u + \alpha_v = 2 = w_M(u, v)$.*

Proof Level 1 gadgets that correspond to variables set to true have blocking edges. More precisely, for every variable X_k set to true, (x_k, y_k) is a blocking edge to M and we have $\alpha_{x_k} = \alpha_{y_k} = 1$. Thus $\alpha_{x_k} + \alpha_{y_k} = 2 = w_M(x_k, y_k)$. Similarly, consider any level 2 or level 3 gadget that is in unstable state: such a gadget has a blocking edge within it, say (p_0^c, q_0^c) or (p_4^c, q_4^c) or (s_3^c, t_3^c) , and both endpoints of such an edge have their α -values set to 1. For the D -gadget, (d_1, d_3) is a blocking edge and we have $\alpha_{d_1} = \alpha_{d_3} = 1$. There are no blocking edges to M in the Z -gadget or in a level 0 gadget. Thus all intra-gadget blocking edges are covered. \square

Claim 5 *Let (u, v) be an intra-gadget edge that is non-blocking. Then $\alpha_u + \alpha_v \geq w_M(u, v)$.*

Proof For any edge (z_i, z_j) where $i, j \in \{0, 1, \dots, 5\}$, we have $\alpha_{z_i} + \alpha_{z_j} = 0 = w_M(z_i, z_j)$. Similarly, all edges within the D -gadget are covered. For any variable X_i set to false: $\alpha_{x_i} + \alpha_{y_i} = 0 = w_M(x_i, y_i)$ and similarly, $\alpha_{x'_i} + \alpha_{y_i} = 0 = w_M(x'_i, y_i)$. For any variable X_k set to true: $\alpha_{x'_k} + \alpha_{y'_k} = -2 = w_M(x'_k, y'_k)$.

For any (a_i^c, b_i^c) , we have $\alpha_{a_i^c} + \alpha_{b_i^c} = 0 = w_M(a_i^c, b_i^c)$. Similarly, $\alpha_{a_{2i-1}^c} + \alpha_{b_{2i}^c} = 0 = w_M(a_{2i-1}^c, b_{2i}^c)$, also $\alpha_{a_{2i}^c} + \alpha_{b_{2i-1}^c} = 0 = w_M(a_{2i}^c, b_{2i-1}^c)$ for all i and c .

We also have for all c : $\alpha_{p_1^c} + \alpha_{q_2^c} = 0 = w_M(p_1^c, q_2^c)$ while $\alpha_{p_2^c} + \alpha_{q_1^c} = -2 = w_M(p_2^c, q_1^c)$ and $\alpha_{p_2^c} + \alpha_{q_2^c} = -2 = w_M(p_2^c, q_2^c)$. It is similar for all other edges within level 2 gadgets and also for edges within level 3 gadgets. Thus it is easy to see that for all intra-gadget non-blocking edges (u, v) , we have $\alpha_u + \alpha_v \geq w_M(u, v)$. \square

Claim 6 *Let (u, v) be any inter-gadget edge. Then $\alpha_u + \alpha_v \geq w_M(u, v)$.*

Proof We show that no inter-gadget edge blocks M . The vertices z_0 and z_1 prefer some neighbors in levels 0, 1, 2 to each other and the α -value of each of these neighbors is either 0 or 1. In particular, $\alpha_{x_i} \geq 0$ and $\alpha_{y_i} \geq 0$, $\alpha_{p_1^c} \geq 0$ and $\alpha_{q_0^c} \geq 0$, and so on while $\alpha_{a_i^c} = \alpha_{b_i^c} = \alpha_{a_i^c} = \alpha_{b_i^c} = 0$ for all i and c . Note that all these vertices prefer their partners in M to z_0 or z_1 ; thus for any such edge e , we have $w_M(e) = 0$. Since $\alpha_{z_i} = 0$ for all i , the edges incident to z_i are covered for all i .

Consider edges between a level 0 vertex and a level 1 vertex, such as (a_1^c, y_j^c) or (b_1^c, x_k^c) : regarding the former edge, we have $w_M(a_1^c, y_j^c) = 0 = \alpha_{a_1^c} + \alpha_{y_j^c}$ and for the latter edge, we have $w_M(b_1^c, x_k^c) = -2 < -1 = \alpha_{b_1^c} + \alpha_{x_k^c}$. It can similarly be verified that every edge between a level 0 vertex and a level 1 vertex is covered.

Consider edges between a level 1 vertex and a level 2 vertex, such as (p_2^c, y_j^c) or (x_k, q_2^c) : recall that (p_0^c, q_2^c) and (x_k, y_k^c) are in M and so $w_M(x_k, q_2^c) = 0$; we set $\alpha_{q_2^c} = -1$ and $\alpha_{x_k} = 1$, thus $\alpha_{x_k} + \alpha_{q_2^c} = w_M(x_k, q_2^c)$. We have $w_M(p_2^c, y_j^c) = -2$ since (p_2^c, q_0^c) and (x_j, y_j^c) are in M and so this edge is covered. It can similarly be verified that every edge between a level 1 vertex and a level 2 vertex is covered.

Consider edges between a level 2 vertex and a level 3 vertex, such as those incident to s_0^c or t_0^c : we have $w_M(s_0^c, q_0^c) = w_M(s_0^c, q_3^c) = 0$ and $\alpha_{s_0^c} = -1$ while $\alpha_{q_0^c} = \alpha_{q_3^c} = 1$. Similarly, $w_M(p_7^c, t_0^c) = w_M(p_4^c, t_0^c) = -2$ and so these edges are covered. It is analogous with edges incident to s_0^c or t_0^c .

Consider any edge e whose one endpoint is in the D -gadget and the other endpoint is outside the D -gadget. It is easy to see that $w_M(e) = -2$, hence this edge is covered. Similarly, inter-gadget edges between levels 0 and 2, levels 0 and 3, and levels 1 and 3 all have weight -2 and hence they are covered. □

Thus we have shown the following theorem.

Theorem 4 *If B has a 1-in-3 satisfying truth assignment then G admits a popular matching.*

Theorem 1 stated in Sect. 1 follows from Theorems 3 and 4. Thus the popular matching problem in a roommates instance on n vertices with complete preference lists is NP-complete for even n .

Acknowledgements Thanks to Chien-Chung Huang for asking us about the complexity of the popular roommates problem with complete preference lists. We are grateful to the reviewers for their helpful comments and suggestions that improved the presentation of the paper.

Funding Open Access funding provided by ELKH Centre for Economic and Regional Studies.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is

not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Biró, P., Irving, R. W., Manlove, D.: Popular matchings in the marriage and roommates problems. In: Proceedings of the 7th International Conference on Algorithms and Complexity (CIAC), pp. 97–108 (2010)
2. Biró, P., McDerimid, E.: Three-sided stable matchings with cyclic preferences. *Algorithmica* **58**(1), 5–18 (2010)
3. Brandl, F., Kavitha, T.: Two problems in max-size popular matchings. *Algorithmica* **81**(7), 2738–2764 (2019)
4. Chung, K.S.: On the existence of stable roommate matchings. *Games Econ Behav* **33**(2), 206–230 (2000)
5. de Condorcet (Marquis de) M-J-A-N: Essai sur l’application de l’analyse à la probabilité des décisions rendues à la pluralité des voix. L’Imprimerie Royale (1785)
6. Condorcet method. https://en.wikipedia.org/wiki/Condorcet_method
7. Cseh, Á.: Popular matchings. In: Endriss U (Ed) Trends in computational social choice, COST (European Cooperation in Science and Technology), pp. 105–122 (2017)
8. Cseh, Á., Kavitha, T.: Popular edges and dominant matchings. *Math. Program.* **172**(1), 209–229 (2018)
9. Cseh, Á., Kavitha, T.: Popular matchings in complete graphs. In: 38th Foundations of Software Technology and Theoretical Computer Science (FSTTCS), pp. 17:1–17:14 (2018)
10. Faenza, Y., Kavitha, T.: Quasi-popular matchings, optimality, and extended formulations. In: 31st ACM-SIAM Symposium on Discrete Algorithms (SODA), 325–344 (2020)
11. Faenza, Y., Kavitha, T., Powers, V., Zhang, X.: Popular matchings and limits to tractability. In: 30th ACM-SIAM Symposium on Discrete Algorithms (SODA), pp. 2790–2809 (2019)
12. Felsenthal, D., Machover, M.: The Measurement of Voting Power. Edward Elgar Publishing, Cheltenham (1998)
13. Gale, D., Shapley, L.S.: College admissions and the stability of marriage. *Am. Math. Mon.* **69**(1), 9–15 (1962)
14. Gärdenfors, P.: Match making: assignments based on bilateral preferences. *Behav. Sci.* **20**, 166–173 (1975)
15. Gupta, S., Misra, P., Saurabh, S., Zehavi, M.: Popular matching in roommates setting is NP-hard. In: 30th ACM-SIAM Symposium on Discrete Algorithms (SODA), pp. 2810–2822 (2019)
16. Huang, C.-C.: Personal communication
17. Huang, C.-C., Kavitha, T.: Popular matchings in the stable marriage problem. *Inf. Comput.* **222**, 180–194 (2013)
18. Huang, C.-C., Kavitha, T.: Near-popular matchings in the roommates problem. *SIAM J. Discrete Math.* **27**(1), 43–62 (2013)
19. Huang, C.-C., Kavitha, T.: Popularity, mixed matchings, and self-duality. In: 28th ACM-SIAM Symposium on Discrete Algorithms (SODA), pp. 2294–2310 (2017)
20. Irving, R.W.: An efficient algorithm for the “stable roommates” problem. *J. Algorithms* **6**, 577–595 (1985)
21. Janssen, J.C.: On even and odd Latin squares. *J. Comb. Theory Ser. A* **69**(1), 173–181 (1995)
22. Kavitha, T., Mestre, J., Nasre, M.: Popular mixed matchings. *Theor. Comput. Sci.* **412**(24), 2679–2690 (2011)
23. Kavitha, T.: A size-popularity tradeoff in the stable marriage problem. *SIAM J. Comput.* **43**(1), 52–71 (2014)
24. Kavitha, T.: Popular half-integral matchings. In: 43rd International Colloquium on Automata, Languages, and Programming (ICALP), pp 22.1–22.13 (2016)
25. Kavitha, T.: Max-size popular matchings and extensions. arXiv preprint [arXiv:1802.07440](https://arxiv.org/abs/1802.07440) (2018)
26. Lam, C. K., Plaxton, C. G.: On the existence of three-dimensional stable matchings with cyclic preferences. In: 12th International Symposium on Algorithmic Game Theory (SAGT), pp. 329–342 (2019)
27. Manlove, D.F.: *Algorithmics of Matching Under Preferences*. World Scientific, Singapore (2013)
28. Schaefer, T.J.: The complexity of satisfiability problems. In: 10th Annual ACM Symposium on Theory of Computing, pp. 216–226 (1978)

29. Subramanian, A.: A new approach to stable matching problems. *SIAM J. Comput.* **23**(4), 671–700 (1994)
30. Tan, J.J.M.: A necessary and sufficient condition for the existence of a complete stable matching. *J. Algorithms* **12**, 154–178 (1991)
31. Teo, C.-P., Sethuraman, J.: The geometry of fractional stable matchings and its applications. *Math. Oper. Res.* **23**(4), 874–891 (1998)
32. Woeginger, G.J.: Core stability in hedonic coalition formation. In: *International Conference on Current Trends in Theory and Practice of Computer Science*, pp. 33–50 (2013)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.